

Robert Nazaryan and Hayk Nazaryan

**Foundation of Armenian Theory
of Special Relativity
In One Physical Dimension
By Pictures**



Yerevan - 2016
Individual Publication

100 Years Inquisition In Science Is Now Over
Armenian Revolution In Science Has Begun!

2007

Crash Course in Armenian Theory of Special Relativity

September, 2016 - Yerevan, Armenia

Robert Nazaryan and Hayk Nazaryan

**Foundation of Armenian Theory of Special Relativity
In One Physical Dimension by Pictures**

Dedicated to the 25-th Anniversary of Independence of Armenia



Yerevan - 2016
Individual Publication

UDC 530.12

Creation of this book - “**Foundation of Armenian Theory of Special Relativity by Pictures**”, became possible by generous donation from my children:

Nazaryan Gor,
Nazaryan Nazan,
Nazaryan Ara and
Nazaryan Hayk.

I am very grateful to all of them.

We consider the publication of this book as Nazaryan family’s contribution to the renaissance of science in Armenia and the whole world.

Nazaryan R., UDC 530.12

Foundation of Armenian Theory of Special Relativity In One Physical Dimension by Pictures

R. Nazaryan, H. Nazaryan - Yerevan, Ind. Pub., 2016, 76 pages:

© Nazaryan Robert and Nazaryan Hayk, 2016

First Armenian publication – June 2013, Armenia, ISBN: 978-1-4675-6080-1

Illustrated Armenian Publication – August 2016, Armenia, ISBN: 978-9939-0-1981-9

Illustrated English Publication – September 2016, Armenia, ISBN: 978-9939-0-1982-6

Contents

01.	Most General Transformations Between Coordinate Systems and Initial State Condition.....	05
02.	Examining the case of inertial systems.....	09
03.	Implementation of the Relativity Postulate.....	16
04.	Definition of the Coefficient g	26
05.	Examining of the movement of Origin Observing Inertial Systems.....	32
06.	Definition of the Coefficient s	42
07.	Derivation of the Armenian Gamma Functions.....	48
08.	Velocity and Acceleration Formulas of the Observed Test Particle.....	54
09.	Foundation of the Armenian Dynamics.....	65

Our scientific and political articles can be found here.

- <https://yerevan.academia.edu/RobertNazaryan>
- https://archive.org/details/@armenian_theory

*If you have the strong urge to accuse somebody for what you read here,
then don't accuse us, read the sentence to mathematics.
We are simply its messengers only.*

Armenian Theory of Special Relativity Is a New and Solid Mathematical Theory, Because it Satisfies the Conditions to be Called a New Theory

- 1) Our created theory is new, because it was created between the years 2007-2012.
- 2) Our created theory does not contradict former legacy theories of physics.
- 3) The former legacy theory of relativity is a very special case of the Armenian Theory of Relativity (when coefficients are given the values $s = 0$ and $g = -1$).
- 4) All formulas derived by Armenian Theory of Relativity, has a **universal character** because they are the exact mathematical representation of the Nature (*Philosophiae naturalis principia mathematica*).

The book “**Armenian Theory of Relativity**” has been registered at USA Copyright Office on the exact date, 21 December 2012, when all speculative people preaching the end of the world and the end of human species. Our scientific articles hold the following copyrights: TXu 001-338-952 / 2007-02-02, TXu 001-843-370 / 2012-12-21, VAu 001-127-428 / 2012-12-29, TXu 001-862-255 / 2013-04-04, TXu 001-913-513 / 2014-06-21, TXu 001-934-901 / 2014-12-21, TX0 008-218-589 / 2016-02-02

Chapter A

*Most General Transformations
Between Coordinate Systems
And Initial State Condition*

Most General Transformation Forms

- *Time-space coordinates transformations between two reference systems*

Direct transformations

$$\begin{cases} t' = t'(t, x, v) \\ x' = x'(t, x, v) \end{cases}$$

and

Inverse transformations

$$\begin{cases} t = t(t', x', v') \\ x = x(t', x', v') \end{cases}$$

Where all t' , x' , t and x quantities are arbitrary functions.

- *Initial state condition*

When $t = t' = t'' = \dots = 0$

Then origins of all coordinate systems coincide each other on the one origin in 0 point

A_01

A_02

Most General Transformation Equations For Time – Space Coordinates Differentials

- *Direct transformation equations*

$$\left\{ \begin{array}{l} dt' = \frac{\partial t'}{\partial t} dt + \frac{\partial t'}{\partial x} dx + \frac{\partial t'}{\partial v} dv \\ dx' = \frac{\partial x'}{\partial t} dt + \frac{\partial x'}{\partial x} dx + \frac{\partial x'}{\partial v} dv \end{array} \right.$$

A_03

- *Inverse transformation equations*

$$\left\{ \begin{array}{l} dt = \frac{\partial t}{\partial t'} dt' + \frac{\partial t}{\partial x'} dx' + \frac{\partial t}{\partial v'} dv' \\ dx = \frac{\partial x}{\partial t'} dt' + \frac{\partial x}{\partial x'} dx' + \frac{\partial x}{\partial v'} dv' \end{array} \right.$$

A_04

Possible Two Cases depending on Characters of the Observing Coordinate Systems

- In the case of arbitrary observing coordinate systems*

$$\left\{ \begin{array}{l} v \neq \text{constant} \\ v' \neq \text{constant} \end{array} \right. \Rightarrow \left\{ \begin{array}{l} dv \neq 0 \\ dv' \neq 0 \end{array} \right.$$

- In the case of inertial observing coordinate systems*

$$\left\{ \begin{array}{l} v = \text{constant} \\ v' = \text{constant} \end{array} \right. \Rightarrow \left\{ \begin{array}{l} dv = 0 \\ dv' = 0 \end{array} \right.$$

A_05

A_06

Chapter B

*Examining the Case of Inertial Systems
When Time – Space Coordinates
are Homogenous but are Not Isotropic*

In the Case of Observing Inertial Systems We Have the Following Conditions and Transformations

- *Relative velocities are constant*

$$\left\{ \begin{array}{l} v = \text{constant} \\ v' = \text{constant} \end{array} \right. \Rightarrow \left\{ \begin{array}{l} dv = 0 \\ dv' = 0 \end{array} \right.$$

- *Coordinates differentials transformations become*

Direct transformation equations

$$\left\{ \begin{array}{l} dt' = \frac{\partial t'}{\partial t} dt + \frac{\partial t'}{\partial x} dx \\ dx' = \frac{\partial x'}{\partial t} dt + \frac{\partial x'}{\partial x} dx \end{array} \right.$$

Inverse transformation equations

$$\left\{ \begin{array}{l} dt = \frac{\partial t}{\partial t'} dt' + \frac{\partial t}{\partial x'} dx' \\ dx = \frac{\partial x}{\partial t'} dt' + \frac{\partial x}{\partial x'} dx' \end{array} \right.$$

and

Making the Following Notations

- In the case of direct transformations of the coordinates differentials*

Definition of beta coefficients

$$\left\{ \begin{array}{l} \frac{\partial t'}{\partial t} = \beta_1(t, x, v) \\ \frac{\partial t'}{\partial x} = \beta_2(t, x, v) \end{array} \right.$$

and

Definition of gamma coefficients

$$\left\{ \begin{array}{l} \frac{\partial x'}{\partial t} = \gamma_1(t, x, v) \\ \frac{\partial x'}{\partial x} = \gamma_2(t, x, v) \end{array} \right.$$

B_03

- In the case of inverse transformations of the coordinates differentials*

Definition of beta coefficients

$$\left\{ \begin{array}{l} \frac{\partial t}{\partial t'} = \beta'_1(t', x', v') \\ \frac{\partial t}{\partial x'} = \beta'_2(t', x', v') \end{array} \right.$$

and

Definition of gamma coefficients

$$\left\{ \begin{array}{l} \frac{\partial x}{\partial t'} = \gamma'_1(t', x', v') \\ \frac{\partial x}{\partial x'} = \gamma'_2(t', x', v') \end{array} \right.$$

B_04

Direct and Inverse Transformation Equations For Time - Space Coordinates Differentials

- *Coordinates differentials direct transformations*

$$\begin{cases} dt' = \beta_1(t, x, v)dt + \beta_2(t, x, v)dx \\ dx' = \gamma_1(t, x, v)dt + \gamma_2(t, x, v)dx \end{cases}$$

- *Coordinates differentials Inverse transformations*

$$\begin{cases} dt = \beta'_1(t', x', v')dt' + \beta'_2(t', x', v')dx' \\ dx = \gamma'_1(t', x', v')dt' + \gamma'_2(t', x', v')dx' \end{cases}$$

B_05

B_06

In the Case of Homogenous Time – Space Beta and Gamma Coefficients Also Need to Satisfy

- In the case of the coordinates direct transformations*

Property of beta coefficients

$$\begin{cases} \beta_1(t, x, v) \equiv \beta_1(v) \\ \beta_2(t, x, v) \equiv \beta_2(v) \end{cases}$$

and

Property of gamma coefficients

$$\begin{cases} \gamma_1(t, x, v) \equiv \gamma_1(v) \\ \gamma_2(t, x, v) \equiv \gamma_2(v) \end{cases}$$

B_07

- In the case of the coordinates inverse transformations*

Property of beta coefficients

$$\begin{cases} \beta'_1(t', x', v') \equiv \beta'_1(v') \\ \beta'_2(t', x', v') \equiv \beta'_2(v') \end{cases}$$

and

Property of gamma coefficients

$$\begin{cases} \gamma'_1(t', x', v') \equiv \gamma'_1(v') \\ \gamma'_2(t', x', v') \equiv \gamma'_2(v') \end{cases}$$

B_08

In the Case of Homogenous Time – Space,
Coordinates Differentials Transformations
Between Two Inertial Systems Become

Direct transformation equations

$$\begin{cases} dt' = \beta_1(v)dt + \beta_2(v)dx \\ dx' = \gamma_1(v)dt + \gamma_2(v)dx \end{cases}$$

and

Inverse transformation equations

$$\begin{cases} dt = \beta'_1(v')dt' + \beta'_2(v')dx' \\ dx = \gamma'_1(v')dt' + \gamma'_2(v')dx' \end{cases}$$

Reminder

Time – Space is only homogenous but not isotropic,
Therefore all derived formulas are asymmetric.

B_09

But in the Case of Homogeneous Time – Space, Transformations Can be Written Also Without Differentials

- *Transformation equations in natural order form*

Direct transformation equations

$$\begin{cases} t' = \beta_1(v)t + \beta_2(v)x \\ x' = \gamma_1(v)t + \gamma_2(v)x \end{cases}$$

and

Inverse transformation equations

$$\begin{cases} t = \beta'_1(v')t' + \beta'_2(v')x' \\ x = \gamma'_1(v')t' + \gamma'_2(v')x' \end{cases}$$

B_10

- *Transformation equations in legacy form*

Direct transformation equations

$$\begin{cases} t' = \beta_1(v)t + \beta_2(v)x \\ x' = \gamma_2(v)x + \gamma_1(v)t \end{cases}$$

and

Inverse transformation equations

$$\begin{cases} t = \beta'_1(v')t' + \beta'_2(v')x' \\ x = \gamma'_2(v')x' + \gamma'_1(v')t' \end{cases}$$

B_11

Chapter C

Implementation of the Relativity Postulate

Special Theory of Relativity Postulates

- *Special theory of relativity postulates*

1. All fundamental physical laws have the same mathematical functional forms in all inertial systems.
2. There exists a universal constant velocity C , which has the same value in all inertial systems.
3. In all inertial systems time and space are homogeneous (Special Relativity).

C_01

- *Because of the relativity (first) postulate, corresponding coefficients of direct and inverse transformations must be the same mathematical functions*

Beta functions identity

$$\begin{cases} \beta'_1(\) \equiv \beta_1(\) \\ \beta'_2(\) \equiv \beta_2(\) \end{cases}$$

and

Gama functions identity

$$\begin{cases} \gamma'_1(\) \equiv \gamma_1(\) \\ \gamma'_2(\) \equiv \gamma_2(\) \end{cases}$$

C_02

Implementation of First Postulate in Transformation Equations

- *Transformation equations in legacy form*

Direct transformation equations

$$\begin{cases} t' = \beta_1(v)t + \beta_2(v)x \\ x' = \gamma_2(v)x + \gamma_1(v)t \end{cases}$$

and

Inverse transformation equations

$$\begin{cases} t = \beta_1(v')t' + \beta_2(v')x' \\ x = \gamma_2(v')x' + \gamma_1(v')t' \end{cases}$$

C_03

- *Transformation equations in natural order form*

Direct transformation equations

$$\begin{cases} t' = \beta_1(v)t + \beta_2(v)x \\ x' = \gamma_1(v)t + \gamma_2(v)x \end{cases}$$

and

Inverse transformation equations

$$\begin{cases} t = \beta_1(v')t' + \beta_2(v')x' \\ x = \gamma_1(v')t' + \gamma_2(v')x' \end{cases}$$

C_04

Measurements of the Beta and Gamma Coefficients

- *Coefficients which don't have measurements*

$$\left\{ \begin{array}{l} \beta_1 \Rightarrow \text{don't have measurement} \\ \gamma_2 \Rightarrow \text{don't have measurement} \end{array} \right.$$

C_05

- *Coefficients which have measurements*

$$\left\{ \begin{array}{l} \beta_2 \Rightarrow \text{have inverse measurement of velocity } \left(\frac{1}{c}\right) \\ \gamma_1 \Rightarrow \text{have measurement of velocity } (c) \end{array} \right.$$

C_06

Coordinates Transformation Equations In the Form System of Linear Equations

- *Transformation equations in legacy form*

Direct transformation equations

$$\begin{cases} \beta_1(v)t + \beta_2(v)x = t' \\ \gamma_2(v)x + \gamma_1(v)t = x' \end{cases}$$

and

Inverse transformation equations

$$\begin{cases} \beta_1(v')t' + \beta_2(v')x' = t \\ \gamma_2(v')x' + \gamma_1(v')t' = x \end{cases}$$

- *Transformation equations in natural order form*

Direct transformation equations

$$\begin{cases} \beta_1(v)t + \beta_2(v)x = t' \\ \gamma_1(v)t + \gamma_2(v)x = x' \end{cases}$$

and

Inverse transformation equations

$$\begin{cases} \beta_1(v')t' + \beta_2(v')x' = t \\ \gamma_1(v')t' + \gamma_2(v')x' = x \end{cases}$$

Determinants of the Systems of Linear Equations

- *Notations for determinants of the transformation equations*

$$\left\{ \begin{array}{l} D(v) = \begin{vmatrix} \beta_1(v) & \beta_2(v) \\ \gamma_1(v) & \gamma_2(v) \end{vmatrix} \\ D(v') = \begin{vmatrix} \beta_1(v') & \beta_2(v') \\ \gamma_1(v') & \gamma_2(v') \end{vmatrix} \end{array} \right.$$

C_09

- *The determinants formula of the transformation equations*

$$\left\{ \begin{array}{l} D(v) = \beta_1(v)\gamma_2(v) - \beta_2(v)\gamma_1(v) \neq 0 \\ D(v') = \beta_1(v')\gamma_2(v') - \beta_2(v')\gamma_1(v') \neq 0 \end{array} \right.$$

C_10

The Solutions of the Systems of Linear Equations

- For coordinates of the K observing system, we get solutions

$$t = \frac{1}{D(v)} \begin{vmatrix} t' & \beta_2(v) \\ x' & \gamma_2(v) \end{vmatrix} \quad \text{and} \quad x = \frac{1}{D(v)} \begin{vmatrix} \beta_1(v) & t' \\ \gamma_1(v) & x' \end{vmatrix}$$

C_11

- For coordinates of the K' observing system, we get solutions

$$t' = \frac{1}{D(v')} \begin{vmatrix} t & \beta_2(v') \\ x & \gamma_2(v') \end{vmatrix} \quad \text{and} \quad x' = \frac{1}{D(v')} \begin{vmatrix} \beta_1(v') & t \\ \gamma_1(v') & x \end{vmatrix}$$

C_12

Comparison of the Transformation Equations

- New received forms of the transformation equations*

<p><u>Direct transformation equations</u></p> $\left\{ \begin{array}{l} t' = \frac{\gamma_2(v')}{D(v')}t - \frac{\beta_2(v')}{D(v')}x \\ x' = \frac{\beta_1(v')}{D(v')}x - \frac{\gamma_1(v')}{D(v')}t \end{array} \right.$	and	<p><u>Inverse transformation equations</u></p> $\left\{ \begin{array}{l} t = \frac{\gamma_2(v)}{D(v)}t' - \frac{\beta_2(v)}{D(v)}x' \\ x = \frac{\beta_1(v)}{D(v)}x' - \frac{\gamma_1(v)}{D(v)}t' \end{array} \right.$
---	-----	--

C_13

- Original transformation equations in the legacy form*

<p><u>Direct transformation equations</u></p> $\left\{ \begin{array}{l} t' = \beta_1(v)t + \beta_2(v)x \\ x' = \gamma_2(v)x + \gamma_1(v)t \end{array} \right.$	and	<p><u>Inverse transformation equations</u></p> $\left\{ \begin{array}{l} t = \beta_1(v')t' + \beta_2(v')x' \\ x = \gamma_2(v')x' + \gamma_1(v')t' \end{array} \right.$
---	-----	--

C_14

Relations Between Coefficients

- From comparison of the direct transformation equations we get the relations

$$\left\{ \begin{array}{l} \beta_1(v) = + \frac{\gamma_2(v')}{D(v')} \\ \beta_2(v) = - \frac{\beta_2(v')}{D(v')} \end{array} \right. \quad \text{and} \quad \left\{ \begin{array}{l} \gamma_2(v) = + \frac{\beta_1(v')}{D(v')} \\ \gamma_1(v) = - \frac{\gamma_1(v')}{D(v')} \end{array} \right.$$

- From comparison of the inverse transformation equations we get the relations

$$\left\{ \begin{array}{l} \beta_1(v') = + \frac{\gamma_2(v)}{D(v)} \\ \beta_2(v') = - \frac{\beta_2(v)}{D(v)} \end{array} \right. \quad \text{and} \quad \left\{ \begin{array}{l} \gamma_2(v') = + \frac{\beta_1(v)}{D(v)} \\ \gamma_1(v') = - \frac{\gamma_1(v)}{D(v)} \end{array} \right.$$

Grouping the Important Relations

- *Two important relations*

$$\begin{cases} D(v)D(v') = 1 \\ \beta_1(v)\beta_1(v') = \gamma_2(v)\gamma_2(v') \end{cases}$$

C_17

- *First Invariant relation, which we denote as ζ_1*

$$\frac{\beta_2(v)}{\gamma_1(v)} = \frac{\beta_2(v')}{\gamma_1(v')} = \zeta_1$$

C_18

Chapter D

Definition of the Coefficient g

Examining First Invariant Relation

- Coefficient ζ_1 must have the following functional arguments

$$\left\{ \begin{array}{l} \frac{\beta_2(v)}{\gamma_1(v)} = \zeta_1(v) \\ \frac{\beta_2(v')}{\gamma_1(v')} = \zeta_1(v') \end{array} \right.$$

D_01

- Therefore, the coefficient ζ_1 must satisfy the following functional equation

$$\zeta_1(v) = \zeta_1(v')$$

D_02

Finding the Most General Solution for Functional Equation

- *First possible solution, which is not the general solution*

If $|v'| = |v| \Rightarrow$ then ζ_1 is an arbitrary even function

- *Second possible solution, which is the most general solution*

If $|v'| \neq |v| \Rightarrow$ then ζ_1 is constant quantity

Examining the Most General Solution

- ζ_1 function must be a constant quantity

$$\zeta_1(v) = \zeta_1(v') = \zeta_1 = \text{constant}$$

D_05

- Therefore, beta and gamma coefficients relations are constant

$$\frac{\beta_2(v)}{\gamma_1(v)} = \frac{\beta_2(v')}{\gamma_1(v')} = \zeta_1 = \text{constant}$$

D_06

Definition of the Coefficient g

- *From the measurements of the beta and gamma coefficients, we get*

$$\zeta_1 = -g \frac{1}{c^2} = \text{constant}$$

- *Therefore, for the beta coefficients we obtain*

$$\begin{cases} \beta_2(v) = -g \frac{1}{c^2} \gamma_1(v) \\ \beta_2(v') = -g \frac{1}{c^2} \gamma_1(v') \end{cases}$$

Time – Space Coordinates Transformation Equations and Transformations Discriminant Formulas

- *Time – space coordinates direct and inverse transformation equations*

Direct transformation equations

$$\begin{cases} t' = \beta_1(v)t - g\frac{1}{c^2}\gamma_1(v)x \\ x' = \gamma_2(v)x + \gamma_1(v)t \end{cases}$$

and

Inverse transformation equations

$$\begin{cases} t = \beta_1(v')t' - g\frac{1}{c^2}\gamma_1(v')x' \\ x = \gamma_2(v')x' + \gamma_1(v')t' \end{cases}$$

D_09

- *Transformations discriminant formulas*

$$\begin{cases} D(v) = \beta_1(v)\gamma_2(v) + g\frac{1}{c^2}[\gamma_1(v)]^2 \neq 0 \\ D(v') = \beta_1(v')\gamma_2(v') + g\frac{1}{c^2}[\gamma_1(v')]^2 \neq 0 \end{cases}$$

D_10

Chapter E

Examining Origins Movement Of the Observing Inertial Systems

Making Two Abstract – Theoretical Experiments

- *Above mentioned abstract – theoretical experiments conditions*

$$\begin{array}{ccc} \text{For origin of } K' & & \text{For origin of } K \\ \left\{ \begin{array}{l} x' = 0 \\ x = vt \end{array} \right. & \text{and} & \left\{ \begin{array}{l} x = 0 \\ x' = v't' \end{array} \right. \end{array}$$

E_01

- *Conditions (E_01) we need to use in the following transformation equations*

$$\begin{array}{ccc} \text{Direct transformation equations} & & \text{Inverse transformation equations} \\ \left\{ \begin{array}{l} t' = \beta_1(v)t - g\frac{1}{c^2}\gamma_1(v)x \\ x' = \gamma_2(v)x + \gamma_1(v)t \end{array} \right. & \text{and} & \left\{ \begin{array}{l} t = \beta_1(v')t' - g\frac{1}{c^2}\gamma_1(v')x' \\ x = \gamma_2(v')x' + \gamma_1(v')t' \end{array} \right. \end{array}$$

E_02

First Abstract – Theoretical Experiment

- *The condition of the first abstract – theoretical experiment*

$$\begin{cases} x' = 0 \\ x = vt \end{cases}$$

- *Above condition used on transformation equations (E_02)*

From direct transformation equations

$$\begin{cases} t' = [\beta_1(v) - g \frac{v}{c^2} \gamma_1(v)] t \\ 0 = [\gamma_2(v)v + \gamma_1(v)] t \end{cases}$$

and

From inverse transformation equations

$$\begin{cases} t = \beta_1(v') t' \\ vt = \gamma_1(v') t' \end{cases}$$

E_03

E_04

Results of the First Experiment

- *The first abstract-theoretical experiment's important formulas*

$$\begin{cases} \gamma_1(v) = -\gamma_2(v)v \\ v = \frac{\gamma_1(v')}{\beta_1(v')} \end{cases}$$

E_05

- *The first abstract-theoretical experiment's beta coefficient formulas*

$$\beta_1(v') = \frac{1}{\beta_1(v) - g \frac{v}{c^2} \gamma_1(v)}$$

E_06

Second Abstract – Theoretical Experiment

- *The condition of the second abstract – theoretical experiment*

$$\begin{cases} x = 0 \\ x' = v't' \end{cases}$$

- *Above condition used on transformation equations (E_02)*

From direct transformation equations

$$\begin{cases} t' = \beta_1(v)t \\ v't' = \gamma_1(v)t \end{cases}$$

and

From inverse transformation equations

$$\begin{cases} t = [\beta_1(v') - g \frac{v'}{c^2} \gamma_1(v')] t' \\ 0 = [\gamma_2(v')v' + \gamma_1(v')] t' \end{cases}$$

E_07

E_08

Results of the Second Experiment

- *The second abstract-theoretical experiment's important formulas*

$$\begin{cases} \gamma_1(v') = -\gamma_2(v')v' \\ v' = \frac{\gamma_1(v)}{\beta_1(v)} \end{cases}$$

E_09

- *The second abstract-theoretical experiment's beta coefficient formulas*

$$\beta_1(v) = \frac{1}{\beta_1(v') - g \frac{v'}{c^2} \gamma_1(v')}$$

E_10

Two Experiments Results Written Together

- *First group of coefficients relations*

$$\left\{ \begin{array}{l} \gamma_1(v) = -\gamma_2(v)v \\ \gamma_1(v') = -\gamma_2(v')v' \end{array} \right. \Rightarrow \left\{ \begin{array}{l} \beta_2(v) = g \frac{v}{c^2} \gamma_2(v) \\ \beta_2(v') = g \frac{v'}{c^2} \gamma_2(v') \end{array} \right.$$

- *Second group of coefficients relations*

$$\left\{ \begin{array}{l} \beta_1(v') = \frac{1}{\beta_1(v) + g \frac{v^2}{c^2} \gamma_2(v)} \\ \beta_1(v) = \frac{1}{\beta_1(v') + g \frac{v'^2}{c^2} \gamma_2(v')} \end{array} \right.$$

Relations Between Relative Velocities

- *Relations between inverse and direct relative velocities*

$$\left\{ \begin{array}{l} v' = \frac{\gamma_1(v)}{\beta_1(v)} \\ v = \frac{\gamma_1(v')}{\beta_1(v')} \end{array} \right. \Rightarrow \left\{ \begin{array}{l} v' = -\frac{\gamma_2(v)}{\beta_1(v)} v \\ v = -\frac{\gamma_2(v')}{\beta_1(v')} v' \end{array} \right.$$

E_13

- *Relative velocities satisfy the property*

$$v'' = -\frac{\gamma_2(v')}{\beta_1(v')} v' = v \Rightarrow v'' \equiv v$$

E_14

Transformations Discriminants Formulas

- *First group of discriminants formulas*

$$\begin{cases} D(v) = \gamma_2(v) \left[\beta_1(v) + g \frac{v^2}{c^2} \gamma_2(v) \right] \neq 0 \\ D(v') = \gamma_2(v') \left[\beta_1(v') + g \frac{v'^2}{c^2} \gamma_2(v') \right] \neq 0 \end{cases}$$

- *Second group of discriminants formulas*

$$\begin{cases} D(v) = \beta_1(v) \gamma_2(v) \left(1 - g \frac{vv'}{c^2} \right) \neq 0 \\ D(v') = \beta_1(v') \gamma_2(v') \left(1 - g \frac{vv'}{c^2} \right) \neq 0 \end{cases}$$

E_15

E_16

Direct and Inverse Transformation Equations

- *First form of transformation equations*

Direct transformation equations

$$\begin{cases} t' = \beta_1(v)t + g\frac{v}{c^2}\gamma_2(v)x \\ x' = \gamma_2(v)(x - vt) \end{cases}$$

and

Inverse transformation equations

$$\begin{cases} t = \beta_1(v')t' + g\frac{v'}{c^2}\gamma_2(v')x' \\ x = \gamma_2(v')(x' - v't') \end{cases}$$

E_17

- *Second form of transformation equations*

Direct transformation equations

$$\begin{cases} t' = \beta_1(v)\left(t - g\frac{v'}{c^2}x\right) \\ x' = \gamma_2(v)(x - vt) \end{cases}$$

and

Inverse transformation equations

$$\begin{cases} t = \beta_1(v')\left(t' - g\frac{v}{c^2}x'\right) \\ x = \gamma_2(v')(x' - v't') \end{cases}$$

E_18

Chapter F

Definition of the Coefficient S

For simplicity purposes we will use the beta and gamma coefficients without index.

$$\begin{cases} \beta_1() \Rightarrow \beta() \\ \gamma_2() \Rightarrow \gamma() \end{cases}$$

We Need to Use the Following Previous Results

- From (C_15) and (C_16) we have the following relations between coefficients

$$\left\{ \begin{array}{l} \beta(v) = + \frac{\gamma(v')}{D(v')} \\ \gamma(v) = + \frac{\beta(v')}{D(v')} \end{array} \right. \quad \text{and} \quad \left\{ \begin{array}{l} \beta(v') = + \frac{\gamma(v)}{D(v)} \\ \gamma(v') = + \frac{\beta(v)}{D(v)} \end{array} \right.$$

F_01

- From (E_13) we have the following relations between relative velocities

$$\left\{ \begin{array}{l} v' = -\frac{\gamma(v)}{\beta(v)}v \\ v = -\frac{\gamma(v')}{\beta(v')}v' \end{array} \right.$$

F_02

Second Invariant Relation

- From (F_01) and (F_02) we get second invariant relation, which we denote as ζ_2

$$\frac{\beta(v') - \gamma(v')}{\gamma(v')v'} = \frac{\beta(v) - \gamma(v)}{\gamma(v)v} = \zeta_2$$

- For the most general solution, the above invariant relation must be constant

$$\frac{\beta(v') - \gamma(v')}{\gamma(v')v'} = \frac{\beta(v) - \gamma(v)}{\gamma(v)v} = \zeta_2 = \text{constant}$$

Definition of the Coefficient S

- *From the measurements of beta and gamma coefficients, we get*

$$\zeta_2 = s \frac{1}{c} = \text{constant}$$

F_05

- *Therefore, we can write the second invariant relation with a new coefficient S*

$$\frac{\beta(v') - \gamma(v')}{\gamma(v')v'} = \frac{\beta(v) - \gamma(v)}{\gamma(v)v} = s \frac{1}{c}$$

F_06

Formulas for Beta Coefficients

F_07

$$\left\{ \begin{array}{l} \beta(v) = \gamma(v) \left(1 + s \frac{v}{c} \right) \\ \beta(v') = \gamma(v') \left(1 + s \frac{v'}{c} \right) \end{array} \right.$$

From this point on, all transformation equations and other important relativistic formulas we will name “Armenian”. This is the best way to distinguish between the legacy and the new theory of relativity and their corresponding relativistic formulas.

Also, this research is the accumulation of practically 50 years of obsessive thinking about the natural laws of the universe. It was done in Armenia by an Armenian and the original manuscripts were written in Armenian. This research is purely from the mind of an Armenian and from the land of Armenia, therefore we can call it by its rightful name.

Formulas of the Armenian Theory of Relativity

- *Armenian relation formula between relative velocities*

$$\left\{ \begin{array}{l} v' = -\frac{v}{1 + s\frac{v}{c}} \\ v = -\frac{v'}{1 + s\frac{v'}{c}} \end{array} \right. \Rightarrow \left(1 + s\frac{v}{c}\right)\left(1 + s\frac{v'}{c}\right) = 1$$

F_08

- *Armenian direct and inverse transformation equations*

Armenian direct transformation equations

$$\left\{ \begin{array}{l} t' = \gamma(v) \left[\left(1 + s\frac{v}{c}\right)t + g\frac{v}{c^2}x \right] \\ x' = \gamma(v)(x - vt) \end{array} \right.$$

Armenian inverse transformation equations

$$\left\{ \begin{array}{l} t = \gamma(v') \left[\left(1 + s\frac{v'}{c}\right)t' + g\frac{v'}{c^2}x' \right] \\ x = \gamma(v')(x' - v't') \end{array} \right.$$

F_09

Chapter G

Derivation of the Armenian Gamma Functions

Armenian Invariant Interval Between Two Events

- *Armenian transformation equations in the same measurement coordinates*

Armenian direct transformation equations

$$\begin{cases} ct' = \gamma(v) \left[\left(1 + s \frac{v}{c}\right) ct + g \frac{v}{c} x \right] \\ x' = \gamma(v) \left(x - \frac{v}{c} ct \right) \end{cases}$$

and

Armenian inverse transformation equations

$$\begin{cases} ct = \gamma(v') \left[\left(1 + s \frac{v'}{c}\right) ct' + g \frac{v'}{c} x' \right] \\ x = \gamma(v') \left(x' - \frac{v'}{c} ct' \right) \end{cases}$$

G_01

- *Quadratic form of the Armenian invariant interval*

$$\mathfrak{G}^2 = (ct')^2 + s(ct')x' + gx'^2 = (ct)^2 + s(ct)x + gx^2$$

G_02

Reciprocal Calculation of the Armenian Interval

- *Reciprocal substitution coordinates into Armenian interval formulas (G_02)*

$$\left\{ \begin{array}{l} \mathfrak{E}^2 = [\gamma(v)]^2 \left(1 + s \frac{v}{c} + g \frac{v^2}{c^2} \right) [(ct)^2 + s(ct)x + gx^2] \\ \mathfrak{E}^2 = [\gamma(v')]^2 \left(1 + s \frac{v'}{c} + g \frac{v'^2}{c^2} \right) [(ct')^2 + s(ct')x' + g(x')^2] \end{array} \right.$$

- *Above Armenian interval expressions must be equal original interval formulas*

$$\left\{ \begin{array}{l} \mathfrak{E}^2 = (ct)^2 + s(ct)x + gx^2 \\ \mathfrak{E}^2 = (ct')^2 + s(ct')x' + g(x')^2 \end{array} \right.$$

Equating Two Different Interval Expressions

- *Gamma function of the Armenian direct transformation*

$$\gamma_z(v) = \frac{1}{\sqrt{1 + s\frac{v}{c} + g\frac{v^2}{c^2}}}$$

G_05

- *Gamma function of the Armenian inverse transformation*

$$\gamma_z(v') = \frac{1}{\sqrt{1 + s\frac{v'}{c} + g\frac{v'^2}{c^2}}}$$

G_06

First Group of Important Formulas

- *Armenian transformation equations discriminants values*

$$\begin{cases} D(v) = [\gamma_z(v)]^2 \left(1 + s \frac{v}{c} + g \frac{v^2}{c^2} \right) = 1 \\ D(v') = [\gamma_z(v')]^2 \left(1 + s \frac{v'}{c} + g \frac{v'^2}{c^2} \right) = 1 \end{cases}$$

- *Armenian gamma functions first group of important relations*

$$\begin{cases} \gamma_z(v') = \gamma_z(v) \left(1 + s \frac{v}{c} \right) \\ \gamma_z(v) = \gamma_z(v') \left(1 + s \frac{v'}{c} \right) \\ \gamma_z(v') v' = -\gamma_z(v) v \end{cases}$$

G_07

G_08

Second Group of Important Formulas

- *This important relation between Armenian gamma functions we use in future for the Armenian energy formulas*

$$\gamma_z(v') \left(1 + \frac{1}{2} s \frac{v'}{c} \right) = \gamma_z(v) \left(1 + \frac{1}{2} s \frac{v}{c} \right)$$

G_09

- *This important relation between Armenian gamma functions we use in future for the Armenian momentum formulas*

$$\gamma_z(v') \left(\frac{1}{2} s + g \frac{v'}{c} \right) + \gamma_z(v) \left(\frac{1}{2} s + g \frac{v}{c} \right) = s \left[\gamma_z(v) \left(1 + \frac{1}{2} s \frac{v}{c} \right) \right]$$

G_10

Chapter H

Velocity and Acceleration Formulas Of the Observed Test Particle

Notations for the Test Particle Velocities and Accelerations

- *Notation for the moving test particle velocities*

$$\begin{cases} u = \frac{dx}{dt} \\ u' = \frac{dx'}{dt'} \end{cases}$$

H_01

- *Notation for the moving test particle accelerations*

$$\begin{cases} b = \frac{du}{dt} = \frac{d^2x}{dt^2} \\ b' = \frac{du'}{dt'} = \frac{d^2x'}{dt'^2} \end{cases}$$

H_02

Time Derivatives of the Armenian Transformation Equations

- *Time derivatives of the Armenian direct transformation equations*

$$\begin{cases} \frac{dt'}{dt} = \gamma_z(v) \left(1 + s \frac{v}{c} + g \frac{vu}{c^2} \right) \\ \frac{dx'}{dt} = \gamma_z(v)(u - v) \end{cases}$$

- *Time derivatives of the Armenian inverse transformation equations*

$$\begin{cases} \frac{dt}{dt'} = \gamma_z(v') \left(1 + s \frac{v'}{c} + g \frac{v'u'}{c^2} \right) \\ \frac{dx}{dt'} = \gamma_z(v')(u' - v') \end{cases}$$

H_03

H_04

Relations of the Time Differentials

- *First form of relations of the time differentials*

$$\begin{cases} \frac{dt'}{dt} = \gamma_z(v) \left(1 + s \frac{v}{c} + g \frac{vu}{c^2} \right) \\ \frac{dt}{dt'} = \gamma_z(v') \left(1 + s \frac{v'}{c} + g \frac{v'u'}{c^2} \right) \end{cases}$$

H_05

- *Second form of relations of the time differentials*

$$\begin{cases} \frac{dt'}{dt} = \gamma_z(v') \left(1 - g \frac{v'u}{c^2} \right) \\ \frac{dt}{dt'} = \gamma_z(v) \left(1 - g \frac{vu'}{c^2} \right) \end{cases}$$

H_06

Moving Test Particle Velocity Formulas

- *Test particle velocity with respect to the inertial system K'*

$$\frac{dx'}{dt'} = u' = \frac{u - v}{1 + s\frac{v}{c} + g\frac{vu}{c^2}}$$

- *Test particle velocity with respect to the inertial system K*

$$\frac{dx}{dt} = u = \frac{u' - v'}{1 + s\frac{v'}{c} + g\frac{v'u'}{c^2}}$$

H_07

H_08

Armenian Addition and Subtraction Formulas for Velocities

- *Armenian addition and subtraction formulas, expressed by direct relative velocity*

$$\left\{ \begin{array}{l} u = u' \oplus v = \frac{\left(1 + s \frac{v}{c}\right) u' + v}{1 - g \frac{vu'}{c^2}} \\ u' = u \ominus v = \frac{u - v}{1 + s \frac{v}{c} + g \frac{vu}{c^2}} \end{array} \right.$$

H_09

- *Armenian addition and subtraction formulas, expressed by inverse relative velocity*

$$\left\{ \begin{array}{l} u' = u \oplus v' = \frac{\left(1 + s \frac{v'}{c}\right) u + v'}{1 - g \frac{v'u}{c^2}} \\ u = u' \ominus v' = \frac{u' - v'}{1 + s \frac{v'}{c} + g \frac{v'u'}{c^2}} \end{array} \right.$$

H_10

Gamma Function Formulas for the Test Particle Moving by Arbitrary Velocity

- *Armenian gamma function formula with respect to the inertial system K*

$$\gamma_z(u) = \frac{1}{\sqrt{1 + s\frac{u}{c} + g\frac{u^2}{c^2}}}$$

- *Armenian gamma function formula with respect to the inertial system K'*

$$\gamma_z(u') = \frac{1}{\sqrt{1 + s\frac{u'}{c} + g\frac{u'^2}{c^2}}}$$

Moving Test Particle Gamma Functions Transformations

- *First form of the gamma functions transformation formulas*

$$\begin{cases} \gamma_z(u) = \gamma_z(v)\gamma_z(u')\left(1 - g\frac{vu'}{c^2}\right) \\ \gamma_z(u') = \gamma_z(v')\gamma_z(u)\left(1 - g\frac{v'u}{c^2}\right) \end{cases}$$

H_13

- *Second form of the gamma functions transformation formulas*

$$\begin{cases} \gamma_z(u) = \gamma_z(v')\gamma_z(u')\left(1 + s\frac{v'}{c} + g\frac{v'u'}{c^2}\right) \\ \gamma_z(u') = \gamma_z(v)\gamma_z(u)\left(1 + s\frac{v}{c} + g\frac{vu}{c^2}\right) \end{cases}$$

H_14

Few More Relations Between Armenian Gamma Functions

- *Test particle gamma functions relation formulas*

$$\left\{ \begin{array}{l} \frac{\gamma_z(u)}{\gamma_z(u')} = \gamma_z(v) \left(1 - g \frac{vu'}{c^2}\right) = \gamma_z(v') \left(1 + s \frac{v'}{c} + g \frac{v'u'}{c^2}\right) \\ \frac{\gamma_z(u')}{\gamma_z(u)} = \gamma_z(v') \left(1 - g \frac{v'u}{c^2}\right) = \gamma_z(v) \left(1 + s \frac{v}{c} + g \frac{vu}{c^2}\right) \end{array} \right.$$

- *From (H_15) we get some interesting relations*

$$\left\{ \begin{array}{l} \sqrt{1 + s \frac{v}{c} + g \frac{v^2}{c^2}} \sqrt{1 + s \frac{v'}{c} + g \frac{v'^2}{c^2}} = \left(1 + s \frac{v}{c} + g \frac{vu}{c^2}\right) \left(1 + s \frac{v'}{c} + g \frac{v'u'}{c^2}\right) \\ \sqrt{1 + s \frac{v}{c} + g \frac{v^2}{c^2}} \sqrt{1 + s \frac{v'}{c} + g \frac{v'^2}{c^2}} = \left(1 - g \frac{vu'}{c^2}\right) \left(1 - g \frac{v'u}{c^2}\right) \end{array} \right.$$

Time Differentials New Relations

- *Time differentials new relations for observed test particle*

$$\left\{ \begin{array}{l} \frac{dt}{dt'} = \frac{\gamma_z(u)}{\gamma_z(u')} \\ \frac{dt'}{dt} = \frac{\gamma_z(u')}{\gamma_z(u)} \end{array} \right. \Rightarrow \frac{dt}{\gamma_z(u)} = \frac{dt'}{\gamma_z(u')} = dt$$

H_17

- *Time differentials new relations for two special cases*

$$\left\{ \begin{array}{l} u' = 0 \\ u = 0 \end{array} \right. \Rightarrow \left\{ \begin{array}{l} \frac{dt}{dt'} = \gamma_z(v) \\ \frac{dt'}{dt} = \gamma_z(v') \end{array} \right.$$

H_18

Moving Test Particle Acceleration Formulas

- *Test particle accelerations transformation formulas*

$$\left\{ \begin{array}{l} b' = \frac{1}{\gamma_z^3(v) \left(1 + s \frac{v}{c} + g \frac{vu}{c^2} \right)^3} b = \frac{1}{\gamma_z^3(v') \left(1 - g \frac{v'u}{c^2} \right)^3} b \\ b = \frac{1}{\gamma_z^3(v) \left(1 - g \frac{vu'}{c^2} \right)^3} b' = \frac{1}{\gamma_z^3(v') \left(1 + s \frac{v'}{c} + g \frac{v'u'}{c^2} \right)^3} b' \end{array} \right.$$

- *Definition of the invariant Armenian acceleration for observed test particle*

$$b_z = \gamma_z^3(u)b = \gamma_z^3(u')b' = \text{invariant}$$

H_19

H_20

Chapter I

Foundation of the Armenian Dynamics

Armenian Lagrangians of Material Test Particle Moving Free or Under Conservative Forces

- *Armenian Lagrangian of the free moving material particle*

$$\mathcal{L}_z(u) = -m_0 c^2 \sqrt{1 + s \frac{u}{c} + g \frac{u^2}{c^2}}$$

- *Armenian Lagrangian of the material particle moving in conservative field*

$$\mathcal{L}_z(u, x) = -m_0 c^2 \sqrt{1 + s \frac{u}{c} + g \frac{u^2}{c^2}} - U(x)$$

Where m_0 is rest mass of the material test particle.

I_01

I_02

Armenian Energy and Armenian Momentum Formulas

- *Armenian energy formula*

$$E_z(u, x) = \frac{1 + \frac{1}{2}s\frac{u}{c}}{\sqrt{1 + s\frac{u}{c} + g\frac{u^2}{c^2}}} m_0 c^2 + U(x)$$

I_03

- *Armenian momentum formula*

$$P_z(u) = -\frac{\frac{1}{2}s + g\frac{u}{c}}{\sqrt{1 + s\frac{u}{c} + g\frac{u^2}{c^2}}} m_0 c$$

I_04

Approximation of the Armenian Energy and Momentum Formulas

- *Definition of Armenian rest mass*

$$m_{\leq 0} = -\left(g - \frac{1}{4}s^2\right)m_0 \geq 0$$

- *First approximation of Armenian energy and Armenian momentum*

$$\begin{cases} E_{\leq}(u, x) \approx m_0 c^2 + \frac{1}{2} m_{\leq 0} u^2 + U(x) \\ P_{\leq}(u) \approx -\frac{1}{2} s m_0 c + m_{\leq 0} u \end{cases}$$

Armenian Energy and Momentum Formulas for Rest Particle

- *Armenian energy and Armenian momentum values for rest particle*

$$\begin{cases} E_z(0, x) = m_0 c^2 + U(x) \\ P_z(0) = -\frac{1}{2} s m_0 c \end{cases}$$

I_07

- *Armenian formula for infinite free energy – hope for human species*

$$P_z(0) = -\frac{1}{2} s m_0 c$$

I_08

Armenian Energy and Armenian Momentum Formulas Observed From Inertial Systems K and K'

- *Armenian energy and momentum formulas with respect to inertial system K*

$$\begin{cases} E_z \equiv E_z(u, x) = \gamma_z(u) \left(1 + \frac{1}{2} s \frac{u}{c} \right) m_0 c^2 + U(x) \\ P_z \equiv P_z(u) = -\gamma_z(u) \left(\frac{1}{2} s + g \frac{u}{c} \right) m_0 c \end{cases}$$

- *Armenian energy and momentum formulas with respect to inertial system K'*

$$\begin{cases} E'_z \equiv E_z(u', x') = \gamma_z(u') \left(1 + \frac{1}{2} s \frac{u'}{c} \right) m_0 c^2 + U(x') \\ P'_z \equiv P_z(u') = -\gamma_z(u') \left(\frac{1}{2} s + g \frac{u'}{c} \right) m_0 c \end{cases}$$

Armenian Energy and Armenian Momentum Direct and Inverse Transformation Equations for Free Particle

- *Armenian energy and momentum direct transformation equations*

$$\begin{cases} E'_z = \gamma_z(v)(E_z - vP_z) \\ P'_z = \gamma_z(v)\left[\left(1 + s\frac{v}{c}\right)P_z + g\frac{v}{c^2}E_z\right] \end{cases}$$

I_11

- *Armenian energy and momentum inverse transformation equations*

$$\begin{cases} E_z = \gamma_z(v')(E'_z - v'P'_z) \\ P_z = \gamma_z(v')\left[\left(1 + s\frac{v'}{c}\right)P'_z + g\frac{v'}{c^2}E'_z\right] \end{cases}$$

I_12

Reciprocal Observation of the Identical Material Particles Resting in Both Inertial Systems

- *Armenian energy and momentum of the particle resting in the system K'*

$$\begin{cases} E_z(v) = \gamma_z(v) \left(1 + \frac{1}{2}s \frac{v}{c}\right) m_0 c^2 \\ P_z(v) = -\gamma_z(v) \left(\frac{1}{2}s + g \frac{v}{c}\right) m_0 c \end{cases}$$

- *Armenian energy and momentum of the particle resting in the system K*

$$\begin{cases} E_z(v') = \gamma_z(v') \left(1 + \frac{1}{2}s \frac{v'}{c}\right) m_0 c^2 \\ P_z(v') = -\gamma_z(v') \left(\frac{1}{2}s + g \frac{v'}{c}\right) m_0 c \end{cases}$$

I_13

I_14

Very Important Formulas

- *Relations between Armenian energy and Armenian momentum quantities for reciprocal observed identical material particles*

$$\begin{cases} E_z(v') & = E_z(v) \\ P_z(v') + P_z(v) & = -sE_z(v) \end{cases}$$

I_15

- *Armenian full energy formulas for free moving particle*

$$\begin{cases} \left(g\frac{1}{c}E_z\right)^2 + s\left(g\frac{1}{c}E_z\right)P_z + gP_z^2 = g\left(g - \frac{1}{4}s^2\right)(m_0c)^2 \geq 0 \\ \left(g\frac{1}{c}E'_z\right)^2 + s\left(g\frac{1}{c}E'_z\right)P'_z + gP_z'^2 = g\left(g - \frac{1}{4}s^2\right)(m_0c)^2 \geq 0 \end{cases}$$

I_16

Force Acting on Material Particle Moving in Conservative Field

- *Armenian force formulas*

$$\left\{ \begin{array}{l} F_z = \frac{dP_z}{dt} = -\left(g - \frac{1}{4}s^2\right)m_0\gamma_z^3(u)b \\ F'_z = \frac{dP'_z}{dt'} = -\left(g - \frac{1}{4}s^2\right)m_0\gamma_z^3(u')b' \end{array} \right.$$

- *Armenian interpretation of Newton's second law*

$$\left\{ \begin{array}{l} F_z = m_{z0}b_z \\ F'_z = m_{z0}b_z \end{array} \right. \Rightarrow F'_z = F_z$$

Conclusions

We showed that the «Armenian Theory of Special Relativity» is full of fine and difficult ideas to understand, which in many cases seems to conflict with our everyday experiences and legacy conceptions. This new crash course book is the simplified version for broad audiences. This book is not just generalizing transformation equations and all relativistic formulas; It is also without limitations and uses a pure mathematical approach to bring forth new revolutionary ideas in the theory of relativity. It also paves the way to build general theory of relativity and finally for the construction of the unified field theory – the ultimate dream of every truth seeking physicist.

Armenian Theory of Relativity is such a mathematically solid and perfect theory that it cannot be wrong. Therefore, our derived transformation equations and all relativistic formulas have the potential to not just replace legacy relativity formulas, but also rewrite all modern physics. Lorentz transformation equations and other relativistic formulas is a very special case of the Armenian Theory of Relativity when we put $s = 0$ and $g = -1$.

The proofs in this book are very brief, therefore with just a little effort, the readers themselves can prove all the provided formulas in detail. You can find the more detailed proofs of the formulas in our main research book «Armenian Theory of Special Relativity», published in Armenia of June 2013.

In this visual book, you will set your eyes on many new and beautiful formulas which the world has never seen before, especially the crown jewel of the Armenian Theory of Relativity - Armenian energy and Armenian momentum formulas, which can change the future of the human species.

The time has come to reincarnate the ether as a universal reference medium which does not contradict relativity theory. Our theory explains all these facts and peacefully brings together followers of absolute ether theory, relativistic ether theory and dark matter theory. We just need to mention that the absolute ether medium has a very complex geometric character, which has never been seen before.

Our Published Articles and Books

- “Armenian Transformation Equations In 3D (Very Special Case)”, 16 pages, February 2007, USA
- “Armenian Theory of Special Relativity in One Dimension”, Book, 96 pages, **Uniprint**, June 2013, Armenia
- “Armenian Theory of Special Relativity Letter”, **IJRSTP**, Volume 1, Issue 1, April 2014, Bangladesh
- “Armenian Theory of Special Relativity Letter”, 4 pages, **Infinite Energy**, Volume 20, Issue 115, May 2014, USA
- “Armenian Theory of Special Relativity Illustrated”, **IJRSTP**, Volume 1, Issue 2, November 2014, Bangladesh
- “Armenian Theory of Relativity Articles (Between Years 2007 - 2014)”, Book, 42 pages, **LAMBERT Academic Publishing**, February 2016, Germany
- “Armenian Theory of Special Relativity Illustrated”, 11 pages, **Infinite Energy**, Volume 21, Issue 126, March 2016, USA
- “Time and Space Reversal Problems in the Armenian Theory of Asymmetric Relativity”, 17 pages, **Infinite Energy**, Volume 22, Issue 127, May 2016, USA

Inspiration Quotes

“To avoid criticism say nothing, do nothing, be nothing.”

Aristotle

“The secret of change is to focus all your energy not on fighting the old, but on building the new.”

Socrates

“All truth passes through three stages.
First, it is ridiculed, second it is violently opposed, and third, it is accepted as self-evident.”

Arthur Schopenhauer


© Nazaryan Robert and Nazaryan Hayk, 2016, UDC 530.12

First Armenian publication – June 2013, Armenia, ISBN: 978-1-4675-6080-1

Illustrated Armenian Publication – August 2016, Armenia, ISBN: 978-9939-0-1981-9

Illustrated English Publication – September 2016, Armenia, ISBN: 978-9939-0-1982-6

Authors Short Biographies



Robert Nazaryan, a grandson of surviving victims of the Armenian Genocide (1915 - 1921), was born on August 7, 1948 in Yerevan, the capital of Armenia. As a senior in high school he won first prize in the national mathematics Olympiad of Armenia in 1966. Then he attended the Physics department at Yerevan State University from 1966 - 1971 and received his MS in Theoretical Physics. 1971 - 1973 he attended Theological Seminary at Etchmiadzin, Armenia and received Bachelor of Theology degree. For seven years (1978 - 1984) he was imprisoned as a political prisoner in the USSR for fighting for the self-determination of Armenia. He has many ideas and unpublished articles in theoretical physics that are waiting his time to be revealed. Right now he is working to finish “**Armenian Theory of Relativity in 3 Physical Dimensions**”. He has three sons, one daughter and six grandchildren.



Hayk Nazaryan was born on May 12, 1989 in Los Angeles, California. He attended Glendale community Collage from 2009 - 2011, then he transferred to California State University Northridge and got his Master of Science degree in physics 2015. He is now teaching as an adjunct instructor at Glendale Community College.